Rate Of A Chemical Reaction

- 1 Which sample of HCl(aq) reacts at the fastest rate with a 1.0-gram sample of iron fi lings?
 - (1) 10. mL of 1 M HCl(aq) at 10.°C
 - (2) 10. mL of 1 M HCl(aq) at 25°C
 - (3) 10. mL of 3 M HCl(aq) at 10.°C
 - (4) 10. mL of 3 M HCl(aq) at 25°C
- 2 A 5.0-gram sample of Fe(s) is to be placed in 100. milliliters of HCl(aq). Which changes will result in the fastest rate of reaction?
 - (1) increasing the surface area of Fe(s) and increasing the concentration of HCl(aq)
 - (2) increasing the surface area of Fe(s) and decreasing the concentration of HCl(aq)
 - (3) decreasing the surface area of Fe(s) and increasing the concentration of HCl(aq)
 - (4) decreasing the surface area of Fe(s) and decreasing the concentration of HCl(aq)
- 3 Which factors have the greatest effect on the rate of a chemical reaction between AgNO₃(aq) and Cu(s)?
 - (1) solution concentration and temperature
 - (2) solution concentration and pressure
 - (3) molar mass and temperature
 - (4) molar mass and pressure

- 4 For a reaction at equilibrium, which change can increase the rates of the forward and reverse reactions?
 - (1) a decrease in the concentration of the reactants
 - (2) a decrease in the surface area of the products
 - (3) an increase in the temperature of the system
 - (4) an increase in the activation energy of the forward reaction
- 5 The effect of a catalyst on a chemical reaction is to provide a new reaction pathway that results in a different
 - (1) potential energy of the products
 - (2) heat of reaction
 - (3) potential energy of the reactants
 - (4) activation energy
- 6 Addition of a catalyst can speed up a reaction by providing an alternate reaction pathway that has a
 - (1) lower activation energy
 - (2) higher activation energy
 - (3) lower heat of reaction
 - (4) higher heat of reaction
- 7 A catalyst increases the rate of a chemical reaction by
 - (1) providing an alternate reaction pathway
 - (2) providing the required heat of reaction
 - (3) increasing the potential energy of the products
 - (4) increasing the activation energy of the reaction

Base your answers to questions 8 on the information below and on your knowledge of chemistry.

The balanced equation below represents the reaction between a 5.0-gram sample of zinc metal and a 0.5 M solution of hydrochloric acid. The reaction takes place in an open test tube at 298 K and 1 atm in a laboratory activity.

$$Zn(s) + 2HCl(aq) \rightarrow H_2(g) + ZnCl_2(aq) + energy$$

8 State one change in reaction conditions, other than adding a catalyst, that will increase the rate of the reaction.

Base your answers to questions 9 on the information below and your knowledge of chemistry.


Methanol can be manufactured by a reaction that is reversible. In the reaction, carbon monoxide gas and hydrogen gas react using a catalyst. The equation below represents this system at equilibrium.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g) + energy$$

9 State the effect on the rates of both the forward and reverse reactions if no catalyst is used in the system.

Base your answers to questions 10 on the information below and on your knowledge of chemistry.


In a laboratory apparatus, a sample of lead(II) oxide reacts with hydrogen gas at high temperature. The products of this reaction are liquid lead and water vapor. As the reaction proceeds, water vapor and excess hydrogen gas leave the glass tube. The diagram and balanced equation below represent this reaction.



10 State one change in reaction conditions, other than adding a catalyst, that would cause the rate of this reaction to increase.

Base your answers to questions 11 on the information below.

The potential energy diagram and balanced equation shown below represent a reaction between solid carbon and hydrogen gas to produce 1 mole of $C_2H_4(g)$ at 101.3 kPa and 298 K.



11 Identify one change in the reaction conditions, other than adding a catalyst, that can increase the rate of this reaction.

Base your answers to questions 12 on the information below.

Calcium reacts with water. This reaction is represented by the balanced equation below. The aqueous product of this reaction can be heated to evaporate the water, leaving a white solid, $Ca(OH)_2(s)$.

$$Ca(s) + 2H_2O(\ell) \rightarrow Ca(OH)_2(aq) + H_2(g)$$

12 State one change in reaction conditions that will increase the rate of the reaction.

Base your answers to questions 13 on the information below and on your knowledge of chemistry.

Automobile catalytic converters use a platinum catalyst to reduce air pollution by changing emissions such as carbon monoxide, CO(g), into carbon dioxide, $CO_2(g)$. The uncatalyzed reaction is represented by the balanced equation below.

$$2CO(g) + O_2(g) \rightarrow 2CO_2(g) + heat$$

13 Compare the activation energy of the catalyzed reaction to the activation energy of the uncatalyzed reaction.

Base your answers to questions 14 on the information below and on your knowledge of chemistry.

Many breads are made by adding yeast to dough, causing the dough to rise. Yeast is a type of microorganism that produces the catalyst zymase, which converts glucose, $C_6H_{12}O_6$, to ethanol and carbon dioxide gas. The balanced equation for this reaction is shown below.

$$C_6H_{12}O_6(aq) \xrightarrow{zymase} 2C_2H_5OH(aq) + 2CO_2(g)$$

14 Describe how the catalyst, zymase, speeds up this reaction.

Base your answers to questions 15 on the information below and on your knowledge of chemistry.

Ethene and hydrogen can react at a faster rate in the presence of the catalyst platinum. The equation below represents a reaction between ethene and hydrogen.

$$H > C = C < H + H - H \longrightarrow H - C - C - H$$

15 Explain, in terms of activation energy, why the catalyzed reaction occurs at a faster rate.

1 4 2 1 3 1 4 3

5 4

6 1

7 1

- 8 Allow 1 credit. Acceptable responses include, but are not limited to:
 - Increase the surface area of the zinc.
 - Increase the temperature of the reaction.
 - Use a more concentrated HCl(aq) solution.
- 9 Allow 1 credit. Acceptable responses include, but are not limited to:
 - Rate of forward reaction: decreases/slower
 - Rate of reverse reaction: decreases/slower
- 10 Allow 1 credit. Acceptable responses include, but are not limited to:
 - Increase the temperature.
 - Increase the concentration of the hydrogen gas in the tube.
 - Grind the metal oxide to increase its surface area.
- 11 Allow 1 credit. Acceptable responses include, but are not limited to:
 - Increase the temperature.
 - Increase the pressure.
 - Increase the concentration of $H_2(g)$.
 - Increase the surface area of the carbon.
- 12 Allow 1 credit. Acceptable responses include, but are not limited to:
 - Increase the temperature of the water.
 - Increase the surface area of Ca(s).
- 13 Allow 1 credit. Acceptable responses include, but are not limited to:
 - The catalyzed reaction has a lower activation energy than the uncatalyzed reaction.
 - The activation energy is higher for the reaction with no catalyst.
- 14 Allow 1 credit. Acceptable responses include, but are not limited to:
 - Zymase is a catalyst that provides an alternative pathway, which requires less energy.
 - decreases the activation energy
 - changes the reaction mechanism
- 15 Allow 1 credit. Acceptable responses include, but are not limited to:
 - The catalyzed reaction pathway has a lower activation energy than the original reaction.
 - Less energy is needed.